

# Giant condylomata (Buschke-Löwenstein tumours): our case load in surgical treatment and review of the current therapies

M. TRIPOLI, A. CORDOVA, F. MAGGI, F. MOSCHELLA

Chirurgia Plastica e Ricostruttiva, Dipartimento di Discipline Chirurgiche ed Oncologiche, Università degli Studi di Palermo (Italy)

**Abstract. – Background:** Buschke-Löwenstein tumour (BLT) or giant condyloma is a verrucous infiltrating lesion, due to a sexually transmitted virus infection, human papilloma virus subtypes 6 and 11. Poor hygiene, promiscuity, chronic irritation and cellular immunocompromised states are often implicated in its genesis. Typical treatment of giant condyloma includes imiquimod cream, podophillin resin, cryotherapy, laser surgery, tangential shave excision with electrocautery.

**Objective:** The authors report their case load in the treatment of giant condyloma and the review of the modern therapies.

**Methods and Materials:** 27 consecutive patients (18 men, nine women) underwent surgery for giant condylomata of perianal region and externa genitalia at the Department of Plastic Surgery of the University of Palermo, from October 2006 to December 2009. All the patients had been treated before with conservative therapies without significant results. We performed the radical excision with split-thickness skin graft in all the patients.

**Results:** No significant complications have occurred in all the cases. The functional and aesthetic outcome were satisfying. No recurrence of disease were noticed in the follow-up.

**Conclusion:** the radical excision with split-thickness skin graft appears to be a successful option of treatment for Buschke-Löwenstein tumours. Compared to other methods it does not necessitate several stages of treatment, moreover it has the advantage of a lower risk of recurrence, it allows a complete histologic examination, the healing process is rapid, the improvement of quality of patients's life is significant.

## Key Words:

Giant condyloma, Buschke-Löwenstein tumour, Surgical excision, Split-thickness skin graft.

## Introduction


Inizialmente descritto da Buschke e Löwenstein<sup>1</sup>, il giant condyloma è una rara manifestazione di condiloma acuminatum, che può superare i 10 a 15 cm di diametro, dovuta a un'infezione virale con il papillomavirus umano dei tipi 6 e 11, trasmessa per contatto sessuale, autoinoculazione, contatto con materiali infetti<sup>2</sup>.

Macroscopicamente appare come una massa polipoidica, cauliflorata, caratterizzata da una crescita lenta, infiltrazione locale, distruzione tissutale contigua, con alta tendenza alla recidiva e alla produzione di fistole o ascessi attorno all'area colpita. Le localizzazioni più frequenti sono la superficie della vulva, lo scroto, il pene, il perineo e la regione perianale (Figura 1), mentre l'implicazione del retto e della vescica è estremamente rara<sup>3</sup>. È controversa l'idea che il tumore di Buschke-Löwenstein debba essere considerato una lesione premaligna con un alto potenziale di degenerazione maligna o una manifestazione di carcinoma verrucoso, poiché alcuni Autori ne riportano la trasformazione in carcinoma squamoso dopo decenni di crescita<sup>4</sup>.

## Modern Therapies

Differenti tipi di trattamento sono riportati per il giant condyloma. Tuttavia, a causa della sua rarità, la letteratura è composta principalmente da casi clinici e manca di studi controllati. Il tipo di trattamento dipende da molti fattori, tra cui le dimensioni, la localizzazione del giant condyloma, le terapie precedenti. Le recidive sono frequenti, specialmente dopo una terapia conservativa.

La soluzione di podofilossolo e il gel sono le prime opzioni di trattamento per il condiloma acuminatum. Con la soluzione al 0.5%, le recidive sono state riscontrate in un



**Figure 1.** Clinical presentation of a Buschke-Löwenstein tumour involving the pubis, the perineum, the left groin and the anorectal region.

third of the patients during the first month after the treatment<sup>5</sup>. Podofilox gel, removes more than 64% of the warts after 8 weeks of treatment<sup>6</sup> with moderate local adverse reaction. It is reported an high risk of recurrence, probably because of the presence of virus in adjacent normal-appearing tissue<sup>7</sup>.

The antimetabolite 5-fluorouracil<sup>8,9</sup>, the bleomycin combined with cisplatin or methotrexate<sup>10,11</sup>, the  $\alpha$ -interferon<sup>12</sup> in topical or systemic administration, are possible option of management to reduce the size of the giant condyloma.

However, there are no guidelines or standard doses, the treatments are expensive and require multiple administrations. In addition, the adverse events limit the therapy.

Imiquimod is an immune-response modifying agent that induces the synthesis of interferon  $\alpha$ , interleukine 1, 6 and 8 and other cytokines. Topical 5% imiquimod cream applied 3 times per week overnight for up to 16 weeks, would completely remove the condylomata acuminata in

37% to 54% of the patients. Frequent adverse reactions are local itching, erythema, skin ulceration. Recurrence occurs in 19% of case after 12 weeks of follow-up<sup>13</sup>.

Cryotherapy by itself or in combination with topical 5-fluorouracil was reported with good results in relatively small Buschke-Löwenstein tumours<sup>14-15</sup>.

Laser: CO<sub>2</sub> or argon laser are reserved for relapsing cases or patients who have not received the previously mentioned treatments as an alternative first-line treatment: its advantages are the hemostatic incision and the wound sterilization<sup>16</sup>.


It is reported the systemic use of  $\alpha$ -interferon combined with neodymium: yttrium-aluminium garnet laser therapy for local recurrence with satisfactory results. However, no large series or long-term results are yet available<sup>17</sup>.

Regional radiotherapy may induce undifferentiation of the tumour cells, stimulating a rapid progression of the lesion to the malignant degeneration<sup>18</sup>.

The surgical excision is still the first line of treatment for Buschke-Löwenstein tumours, with a higher success rate (63%-91%) and lower risk of recurrence<sup>19</sup>. It allows the complete histopathologic examination of the tumour, permitting to exclude squamous cell carcinoma for which a radical aggressive surgery is mandatory. The Authors shows that the wide excision of the lesion with split-thickness skin graft can provide a good outcome for patients with Buschke-Löwenstein tumours, especially for cases refractory to all forms of medical therapy.

## Patients and Methods

Twenty-seven consecutive patients (18 men, nine women) underwent surgery for giant condylomata of perianal region and external genitalia at the Department of Plastic Surgery of University of Palermo, from October 2006 to December 2009. Mean age was 49 years (range 27-71). Tumours' secretion culture showed *Escherichia coli* and negative-coagulase *Staphylococci sp.* Ten patients have been treated before with podofilox gel, nine with cryotherapy, two with tangential excision by electrocautery, but all them were refractory to the treatments. After local or general anesthesia, the radical excision with 20 mm of margins was performed then, the split-thickness skin graft. The donor site was for all the patients



**Figure 2.** **A**, Preoperative view. **B**, The giant condylomata are excised radically. **C**, Giant condylomata. **D**, Intraoperative view after the split-thickness skin graft. **E**, Post-operative view after seven days.

the medial region of the thigh: the skin graft was thus processed through the skin mesher to expand its size (Figure 2 a-d).

The verrucous lesions localized inside the rectum were excised with CO<sub>2</sub> laser. Antibiotic prophylaxis (2 g of intravenous cefazolin 30 minutes before the operation) was administrated in all the patients.


## Results

The donor sites healed within 7-14 days in all the cases. The attachment of the skin graft was complete in 23 patients (Figure 2e). In four cas-

es, after the *Pseudomonas aeruginosa* infection, a partial graft failure occurred, resulting in a prolonged healing process by second intention, with frequent medications (three times a week) and post-operative antibiotic administration for a mean of seven days.

The lymphoedema was present in three patients at the genitalia, they were treated with manual lymph drainage with a complete resolution after four weeks. In one patient the shortage of the graft at the groin determined subacute pain which required physical therapy for three months and non-steroidal anti-inflammatory drugs (NSAIDs) for two weeks, with complete resolution of the symptoms.





**Figure 3.** Post-operative view after. **A**, Two months. **B**, Six months. **C**, One year.

The histopathologic analysis reported for all the cases giant condylomata with intraepithelial hyperplasia without cellular atypia. It was not possible to determine which viral subtypes were involved. All the patients were visited in the Institute during the follow-up (range 36-48 months).

No recurrences were noticed in all of them (Figure 3a-c). All the patients compiled a form about their satisfactory for functional and cosmetic results. The 94% of the patients considered excellent the final outcome.

## Discussion

Buschke-Löwenstein tumour or giant condyloma origins from condylomata acuminata, due to the infection by the *human papillomavirus*. An estimated rate of 30% to 50% of sexually active adults has genital human papillomavirus, and 2% presents genital condylomata acuminata<sup>20</sup>. The giant condyloma differs from condylomata acuminata because it can invade and destroy the adjacent

tissue. It is extremely rare, so experience with the treatment of these tumours is difficult to accumulate. Its histological pattern is similar to the verrucous carcinoma, thus many Authors accept to include both the lesions under the term Buschke-Löwenstein tumours<sup>18</sup>. It has been reported malignant transformation in squamous cell carcinoma in 30-50% of patients<sup>21</sup>. The coexistence of Buschke-Löwenstein tumour and squamous cell carcinoma in the same site, changes the behaviour and progress of the lesion<sup>19</sup>. For this reason we believe that the only consistently effective therapy is wide surgical excision of tumour with 20 mm margins, followed by the split-thickness skin graft. This procedure is the best treatment of choice for giant condylomata, permitting a complete histopathologic examination, a more rapid healing and improved final results, with a lower risk of recurrences. Alternative treatment to surgery as laser surgery, photodynamic therapy, cryotherapy, may also be applied in cases of rectal or vulvar localization, only after a bioptic histologic analysis to exclude the present of a squamous cell carcinoma.

## References

- 1) BUSCHKE A, AND LÖWENSTEIN L. Über Carcinomähnliche Condylomata acuminata de Penis. Klin Wochenschr 1925; 4: 1726-1728.
- 2) AMBRIZ-GONZÁLEZ G, ESCOBEDO-ZAVALA LC, CARILLO DE LA MORA F, ORTIZ-ARRIAGA A, CORDERO-ZAMORA A, CORONA-NAKAMURA A, LÓPEZ RAMÍREZ MK, VELÁZQUEZ RAMÍREZ GA. Buschke-Löwenstein tumour in childhood. A case report. J Pediatr Surg 2005; 40: 25-27.
- 3) WIEDEMANN A, DIEKMANN WP, HOLTMANN G, KRACHT H. Report of a case of giant condyloma (Buschke-Löwenstein tumour) localized in the bladder. J Urol 1995; 153: 1222-1224.
- 4) NEMESIO C, MIROWSKI GW, CHUANG TY. Human papillomavirus: clinical significance and malignant potential. Int J Dermatol 2001; 40: 373-379.
- 5) KIRBY P, DUNNE A, KING DH, COREY L. Double-blind randomized clinical trials of self-administered podofilox solution versus vehicle in the treatment of genital warts. Am J Med 1990; 88: 465-469.
- 6) TYRING SK, EDWARDS L, CHERRY LC. Safety and efficacy of 0.5% podofilox gel in the treatment of anogenital warts. Arch Dermatol 1998; 134: 33-38.
- 7) SOBRADO CW, MESTER M, NADALIN W, NAHAS SC, BOCHINI SF, HABR-GAMA A. Radiation-induced total regression of a highly recurrent giant perianal condyloma: report of a case. Dis Colon Rectum 2000; 43: 257-260.
- 8) KREBS H-B. Treatment of genital condylomata with topical 5-fluorouracil. Dermatol Clin 1991; 9: 333-341.
- 9) SWINEHART J, SKINNER R, MCCARTY JM, MILLER HB, TYRING SK, KOREY A, ORENBURG EK. Development of intralesional therapy with fluorouracil/adrenaline injectable gel for management of condyloma acuminata: two phases II clinical studies. Genitourin Med 1997; 73: 481-487.
- 10) BUECKER JW, HEATON CL. Bleomycin sulphate: post-operative intralesional usage for giant condylomata. J Dermatol Surg Oncol 1985; 11: 972-973.
- 11) ILKAY AK, CHODAK GW, VOGELZANG NJ, GERBER GS. Buschke-Löwenstein tumour: therapeutic option including systemic chemotherapy. Urology 1993; 42: 599-602.
- 12) FRIEDMAN-KEIN AE, EVAN LT, CONAN M. Natural interferon alfa for treatment of condyloma acuminata. JAMA 1988; 259: 533-538.
- 13) TYRING S, ARANY I, STANLEY M, TOMAI MA, MILLER RL, SMITH MH, McDERMOTT DJ, SLADE HB. A randomized controlled, molecular study of condylomata acuminata clearance during treatment with imiquimod. J Infect Dis 1998; 178: 551-555.
- 14) CARSON TE. Verrucous carcinoma of the penis: successful treatment with cryosurgery and topical 5-fluorouracil therapy. Arch Dermatol 1978; 114: 1546-1547.
- 15) HUGHES PSH. Cryosurgery of verrucous carcinoma of the penis (Buschke-Löwenstein tumour). Cutis 1979; 24: 43-45.
- 16) APFELBERG DB, MASER MR, LASH H, DRUKER D. CO<sub>2</sub> laser resection for giant perineal condyloma and verrucous carcinoma. Ann Plast Surg 1983; 11: 417-422.
- 17) GILBERT P, BECKERT R. Combination therapy for penile giant Buschke-Löwenstein condyloma. Urol Int 1990; 45: 122-124.
- 18) HATZCHRISTOU DG, APOSTOLIDIS A, TZORTZIS V, HATZIMOURATIDIS K, IOANNIDES E, YANNAKOYORGOS K. An alternative surgical treatment for Buschke-Löwenstein tumours of the penis. Urology 2001; 57: 966-969.
- 19) CHU DO, VEZERIDIS MP, LIBBEY NP, WANEBO HJ. Giant condyloma acuminatum (Buschke-Löwenstein tumour) of the anorectal and perianal regions. Analysis of 42 cases. Dis Colon Rectum 1994; 37: 950-957.
- 20) SEVERSON JL, TYRING SK. Viral disease update. Curr Probl Dermatol 1999; 11: 41-62.
- 21) CREASMAN C, HAAS PA, FOX JR TA, BALAZS M. Malignant transformation of anorectal giant condyloma acuminatum (Buschke-Löwenstein tumour). Dis Colon Rectum 1989; 32: 481-487.